arXiv:1208.2706v1 [quant-ph] 13 Aug 2012

Genuinely Multipartite Concurrence of N-qubit X-matrices

S. M. Hashemi Rafsanjani,[1,*] M. Huber,[2] C. J. Broadbent,[1] and J. H. Eberly[1]
[1] *Rochester Theory Center and the Department of Physics & Astronomy, University of Rochester, Rochester, New York 14627*
[2] *University of Bristol, Department of Mathematics, BS8 1TW, U.K.*

(Dated: August 15, 2012)

We find an algebraic formula for the N-partite concurrence of N qubits in an X-matrix. X-matricies are density matrices whose only non-zero elements are diagonal or anti-diagonal when written in an orthonormal basis. We use our formula to study the dynamics of the N-partite entanglement of N remote qubits in generalized N-party Greenberger-Horne-Zeilinger (GHZ) states. We study the case when each qubit interacts with a partner harmonic oscillator. It is shown that only one type of GHZ state is prone to entanglement sudden death; for the rest, N-partite entanglement dies out momentarily. Algebraic formulas for the entanglement dynamics are given in both cases.

I. INTRODUCTION

Even though entanglement was promoted by Schrödinger [1] as a fundamental aspect of quantum theory, and in mathematics it predates quantum mechanics by decades [2], its value as a resource for a wide range of potential applications was not appreciated until recently [3, 4]. Furthermore, although its importance is now largely recognized, witnessing and quantifying the entanglement of arbitrary mixed states are still open questions. In only few-party cases do prescriptions exist for determining the entanglement of a mixed state [5–7].

The problem becomes much more difficult for multipartite entanglement, entanglement shared between more than two parties. Multipartite entanglement is thought to play an essential role in many phenomena including quantum metrology [8] and quantum phase transitions [9]. Previous studies of the dynamics of multipartite entanglement have utilized measures that fail to capture exactly when multipartite entanglement disappears [10–13]. Additionally there have been many attempts to solve the problem of determining the multipartite entanglement of a given state [14–16] (see also references in the paper by Ma *et al.* [17]), but an algebraic and/or numerically efficient prescription has not yet emerged. An algebraic prescription would be especially desirable since it can potentially open the door for a wide range of analytical investigations of entanglement dynamics.

Recently, based on previous works by Pope and Milburn [18] and Love *et al.* [19], a new measure of multipartite entanglement, called genuinely multipartite (GM) concurrence, has been proposed, and it has been shown that GM concurrence is an entanglement monotone [17]. This measure reduces to the I-concurrence [20] for a bipartite system, and to Wootters's original concurrence [5] for two qubits. Additionally, an algebraic formula for a lower bound of the GM concurrence has been found by Ma *et al.* [17]. The lower bound, when calculated for a two-qubit X-matrix, matches the value of Wootters's concurrence.

The X-matrix of Yu and Eberly [21] is a density matrix of N qubits, written in an orthonormal product basis, whose non-zero elements are either diagonal or anti-diagonal. The concurrence of a two-qubit X-matrix takes a very simple form [21] and that is why these two-qubit states have been extensively used in studying the dynamics of entanglement between two qubits in many scenarios [21–24].

Restricted forms of X-matrices of more than two qubits have been used in some recent studies of the dynamics of multipartite entanglement. The entanglement measures utilized in these studies yield qualitative information about the multipartite entanglement [25, 26]. Only for Greenberger-Horne-Zeilinger (GHZ) states that undergo pure dephasing, has there been a successful attempt that uses a geometric measure to give the exact dynamics of N-partite entanglement [27].

In view of the fact that the GM concurrence lower bound, derived in Ref. [17], matches the exact value of concurrence for a two-qubit X-matrix, one might wonder if the lower bound might also be exact for more than a two-qubit X-matrix. In this paper we will prove that this conjecture is correct. The lower bound provided by Ma *et al.* [17] is realized by X-matrices. We thus present an *algebraic* formula for the GM concurrence of an N-qubit X-matrix. This is our principal result and it enables analytic formulation of dynamics of N-partite entanglement in different scenarios.

We will use our result to resolve the dynamics of N-partite entanglement of N-qubit GHZ states, which are superpositions of two bit-flipped basis states, when each qubit (two-level atom) interacts locally with an additional system. We assume the additional systems are single-mode cavities, initially in the vacuum state [22, 25]. We will identify two kinds of GHZ states, based on whether or not they experience entanglement sudden death (ESD) [23]. We will show that most of the GHZ states do not experience ESD and only the GHZ state for which, initially, the two-level atoms are all excited or all in their ground states experiences ESD. In each case,

---

[*] hashemi@pas.rochester.edu

we will present algebraic formulas giving the evolution of the GM concurrence.

II. GENUINELY MULTIPARTITE CONCURRENCE

In this section we give a brief introduction to N-partite entanglement and GM concurrence.

N-partite entanglement is defined by its opposite, biseparability. A pure N-partite system $|\psi\rangle \in \mathcal{H}_1 \otimes \mathcal{H}_2 \cdots \otimes \mathcal{H}_N$ is biseparable if there is a bipartition of the N parties $\mathcal{H}_1 \otimes \mathcal{H}_2 \cdots \otimes \mathcal{H}_N = \mathcal{H}_A \otimes \mathcal{H}_B$, where $\mathcal{H}_A = \mathcal{H}_{j_1} \otimes \mathcal{H}_{j_2} \otimes \cdots \otimes \mathcal{H}_{j_k}$, $\mathcal{H}_B = \mathcal{H}_{j_{k+1}} \otimes \mathcal{H}_{j_{k+2}} \otimes \cdots \otimes \mathcal{H}_{j_N}$ for which $|\psi\rangle = |\psi_A\rangle \otimes |\psi_B\rangle$, and $|\psi_A\rangle \in \mathcal{H}_A$ and $|\psi_B\rangle \in \mathcal{H}_B$ [3, 28]. In other words, a pure state is biseparable if it has at least one pure marginal (reduced density matrix). An N-partite state that cannot be written as an ensemble of biseparable states is an N-partite entangled state.

Before introducing GM concurrence, let us introduce the set of all bipartitions of N parties. Each bipartition is a division of the set $\{1, 2, \ldots, N\}$ into two non-overlapping and non-empty subsets. We call the set of all such bipartitions $J = \{J_1, J_2, \ldots J_{2^{N-1}-1}\}$. For example for $N = 3$

$$\begin{aligned} &J_1 = \{1|2,3\}, \quad J_2 = \{2|1,3\}, \quad J_3 = \{3|2,1\}, \\ &J = \{J_1, J_2, J_3\}. \end{aligned} \tag{1}$$

Now if we have a pure state $|\psi\rangle$, to each of the elements of J we can associate two reduced density matrices $\hat{A}(|\psi\rangle, J_j)$, and $\hat{B}(|\psi\rangle, J_j)$ by tracing out either of the subsystems associated with that bipartition. The biseparability of a pure state can be determined by whether for any of the elements of J, $\hat{A}(|\psi\rangle, J_j)$ and $\hat{B}(|\psi\rangle, J_j)$ are pure. If so, $|\psi\rangle$ is biseparable. Thus the purity of the j^{th} bipartition, to be denoted by $\Pi_j(|\psi\rangle)$, is a key parameter.

For a pure state genuinely multipartite concurrence (GM concurrence) is then defined [17] as below.

$$C_{GM}(|\psi\rangle) := \min_j \sqrt{2}\sqrt{1 - \Pi_j(|\psi\rangle)}$$

Clearly, $C_{GM}(|\psi\rangle) \geq 0$ and it is equal to zero if and only if $|\psi\rangle$ is a biseparable state. For a bipartite system this definition reduces to the I-concurrence [20].

To determine whether a mixed state $\hat{\rho}$ is biseparable or not one has to determine whether $\hat{\rho}$ can be written as a convex sum of pure biseparable states. Thus one has to check all the ways $\hat{\rho}$ can be written as a convex sum of pure states (all pure state decompositions). Let us distinguish different pure state decompositions of $\hat{\rho}$ by assigning a superscript, α, to each decomposition

$$\hat{\rho} = \sum_i P_i^\alpha |\psi_i^\alpha\rangle\langle\psi_i^\alpha|. \tag{2}$$

To determine whether a particular pure state decomposition is a sum of biseparable states or not, we can calculate the average pure state GM concurrence for that particular α

$$\begin{aligned} C_\alpha(\hat{\rho}) &= \sum_i P_i^\alpha C_{GM}(|\psi_i^\alpha\rangle) \\ &= \sum_i P_i^\alpha \left\{ \min_j \sqrt{2}\sqrt{1 - \Pi_j(|\psi_i^\alpha\rangle)} \right\}. \end{aligned} \tag{3}$$

Now we are ready to extend the definition of GM concurrence to all mixed states:

$$C_{GM}(\hat{\rho}) = \min_\alpha \; C_\alpha(\hat{\rho}). \tag{4}$$

If $C_{GM}(\hat{\rho}) = 0$ this means that there is an α for which $C_\alpha(\hat{\rho}) = 0$. Then $\hat{\rho}$ can be written as a sum of pure biseparable states, so $\hat{\rho}$ is biseparable. If $C_{GM}(\hat{\rho}) > 0$, there is no α for which the $|\psi_i^\alpha\rangle$'s are all biseparable and thus $\hat{\rho}$ is an N-partite entangled state. The GM concurrence of N parties, as defined in Ref. [17], is a monotone of genuinely multipartite entanglement; it distinguishes between biseparable and N-partite entangled states, is convex, invariant under local unitary transformations, and non-increasing under local operations and classical communications (LOCC) [17].

In [17], Ma *et al.* presented a lower bound for the GM concurrence. Now, we examine this lower bound for an X-matrix of N qubits. If the orthonormal basis for the X-matrix is $\{|0,0,\ldots,0\rangle, |0,0,\ldots,1\rangle, \ldots, |1,1,\ldots,1\rangle\}$, then one can always write an X-matrix in the form given below

$$\hat{X} = \begin{pmatrix} a_1 & & & & & & & z_1 \\ & a_2 & & & & & z_2 & \\ & & \ddots & & & \iddots & & \\ & & & a_n & z_n & & & \\ & & & z_n^* & b_n & & & \\ & & \iddots & & & \ddots & & \\ & z_2^* & & & & & b_2 & \\ z_1^* & & & & & & & b_1 \end{pmatrix}, \tag{5}$$

where $n = 2^{N-1}$, and we require $|z_i| \leq \sqrt{a_i b_i}$ and $\sum_i (a_i + b_i) = 1$ to ensure that $\hat{X}$ is positive and normalized. One can see that the matrix looks like an X, explaining why density matrices in this class are called X-matrices. Without loss of generality we can assume that $\sqrt{a_1 b_1} \geq \sqrt{a_i b_i}$. The lower bound of GM concurrence, derived in [17], for an X-matrix is given by

$$C_{GM} \geq 2\max\{0, |z_i| - w_i\}, \; i = 0, 1, \ldots, n \tag{6}$$

where $w_i = \sum_{j \neq i}^n \sqrt{a_j b_j}$. In the next section we will show that this lower bound is exact for all X-matrices.

III. CONCURRENCE OF N-QUBIT X-STATES

In this section we find an algebraic formula for the GM concurrence of an N-qubit X-matrix.

Before going forward let us introduce some notation. Since we are working with qubits, we can represent each vector (ket) of the above basis as a number from 0 to 2^N-1 written in the binary basis. For example, $|0,0,\cdots,0\rangle = |0\rangle$, $|0,\cdots,0,1\rangle = |1\rangle$, $|0,\cdots,1,0\rangle = |2\rangle$, and so on $|1,\cdots,1,1\rangle = |2^N-1\rangle$. We also denote the bit-flipped states in the same way, $|\bar{i}\rangle = |2^N-i-1\rangle$, e.g., $|\bar{0}\rangle = |2^N-1\rangle$. In places where we need to label the individual qubits, we will do so by using a subscript on the bits,

$$|0_1,0_2,0_3,1_5\rangle = |0\rangle_1 \otimes |0\rangle_2 \otimes |0\rangle_3 \otimes |1\rangle_5. \tag{7}$$

Since we have assumed that $\sqrt{a_1 b_1} \geq \sqrt{a_i b_i}$, it is easy to show that $|z_i| - w_i \leq 0$ for $i > 1$, so that Eq. 6 reduces to

$$C_{GM}(X) \geq 2\max\{0, |z_1| - w_1\} \tag{8}$$

We will show that this bound is actually an equality. First let us prove a lemma that we will utilize in our proof.

Lemma 1. *The GM concurrence of an X-matrix for which* $a_1 b_1 \geq a_i b_i$*, and* $a_j = b_j = 0$ *for all* $j \neq \{i,1\}$*, is*

$$C_{GM}(\hat{X}_{1i}) = 2\max\{0, |z_1| - \sqrt{a_i b_i}\} \tag{9}$$

Proof. We already know that this quantity is a lower bound of GM concurrence. Thus we only need to show that it is also an upper bound. We will do this by mapping $\hat{X_{1i}}$ to a two-qubit density matrix, $\hat{R}$, and then show that $C_{GM}(\hat{X}_{1i})$ is bounded from above by Wootters's concurrence of $\hat{R}$, where $C(\hat{R}) = 2\max\{0, |z_1| - \sqrt{a_i b_i}\}$.

We perform the mapping by focusing on a specific bipartition of the qubits. The four non-zero diagonal elements of $\hat{X}_{1i}$ are $\{a_1, a_i, b_1, b_i\}$, corresponding to projectors $\{|0\rangle\langle 0|, |i-1\rangle\langle i-1|, |\bar{0}\rangle\langle\bar{0}|$, and $|\overline{i-1}\rangle\langle\overline{i-1}|\}$ respectively. Those qubits that contribute 1 to the ket $|i-1\rangle$ we designate as party F. The rest of the qubits we denote as party G. For example, with 7 qubits, which we denote as $(q_1,q_2,q_3,q_4,q_5,q_6,q_7)$, where $i=6$, the basis states are

$$\begin{aligned}
|0\rangle =& |0_1,0_2,0_3,0_4,0_5,0_6,0_7\rangle,\\
|5\rangle =& |0_1,0_2,0_3,0_4,1_5,0_6,1_7\rangle,\\
|127\rangle =& |1_1,1_2,1_3,1_4,1_5,1_6,1_7\rangle,\\
|122\rangle =& |1_1,1_2,1_3,1_4,0_5,1_6,0_7\rangle.
\end{aligned} \tag{10}$$

Then party F is given by qubits (q_1,q_2,q_3,q_4,q_6), and the remaining two qubits, (q_5,q_7), make party G. Under this bipartition, we can write $\hat{X}_{1i}$ using the following basis states,

$$\begin{aligned}
&|{\downarrow_F}\rangle = |0_1,0_2,0_3,0_4,0_6\rangle, && |{\uparrow_F}\rangle = |1_1,1_2,1_3,1_4,1_6\rangle,\\
&|{\downarrow_G}\rangle = \{|0_5,0_7\rangle, && |{\uparrow_G}\rangle = |1_5,1_7\rangle,
\end{aligned}$$

$$\begin{aligned}
\hat{X}_{1i} = &\, a_1|{\downarrow_F\downarrow_G}\rangle\langle{\downarrow_F\downarrow_G}| + b_1|{\uparrow_F\uparrow_G}\rangle\langle{\uparrow_F\uparrow_G}|\\
&+ a_i|{\downarrow_F\uparrow_G}\rangle\langle{\downarrow_F\uparrow_G}| + b_i|{\uparrow_F\downarrow_G}\rangle\langle{\uparrow_F\downarrow_G}|\\
&+ z_1|{\downarrow_F\downarrow_G}\rangle\langle{\uparrow_F\uparrow_G}| + z_1^*|{\uparrow_F\uparrow_G}\rangle\langle{\downarrow_F\downarrow_G}|\\
&+ z_i|{\downarrow_F\uparrow_G}\rangle\langle{\uparrow_F\downarrow_G}| + z_i^*|{\uparrow_F\downarrow_G}\rangle\langle{\downarrow_F\uparrow_G}|,
\end{aligned} \tag{11}$$

We see that if we restrict attention to the subspace defined by the non-zero elements of $\hat{X}_{1i}$ we can map $\hat{X}_{1i}$ to a two qubit density matrix, $\hat{R}$, which, in the basis $\{|{\downarrow_F\downarrow_G}\rangle, |{\downarrow_F\uparrow_G}\rangle$
$, |{\uparrow_F\downarrow_G}\rangle, |{\uparrow_F\uparrow_G}\rangle\}$, reads

$$\hat{X}_{1i} \longrightarrow \hat{R} = \begin{pmatrix} a_1 & & & z_1\\ & a_i & z_i & \\ & z_i^* & b_i & \\ z_1^* & & & b_1 \end{pmatrix}. \tag{12}$$

Now that we have a two-qubit density matrix, we can take advantage of Wootters's concurrence. Note that from each pure-state decomposition (PSD) of $\hat{R}$ one can make a PSD of $\hat{X}_{1i}$ by mapping the basis states of the two-qubit system back to the multi-qubit basis states. We pick the PSD whose average concurrence is the minimum amongst all possible PSD's of $\hat{R}$. Thus

$$\hat{R} = \sum_i p_i|\psi_i\rangle\langle\psi_i|, \quad C(\hat{R}) = \sum_i p_i C(|\psi_i\rangle). \tag{13}$$

In Eq. 13, $C(\hat{R})$ is Wootters's concurrence, which, by definition, is equal to the minimum average concurrence over all possible PSD's of $\hat{R}$. As mentioned before, each pure state $|\psi_i\rangle$ can be mapped back to an N-qubit state ($|\psi_i\rangle \rightarrow |\Psi_i\rangle$), producing a PSD for $\hat{X}_{1i}$,

$$\hat{X}_{1i} = \sum_i p_i|\Psi_i\rangle\langle\Psi_i|. \tag{14}$$

Since GM concurrence is convex by definition, we have

$$C_{GM}(\hat{X}_{1i}) \leq \sum_i p_i C_{GM}(|\Psi_i\rangle). \tag{15}$$

For a pure state the GM concurrence is defined by

$$C_{GM}(|\Psi_i\rangle) = \min_j \sqrt{2}\sqrt{1 - \Pi_j(|\Psi_i\rangle)}, \tag{16}$$

where the minimum is taken over all bipartitions, J, of the N qubits. Therefore, the GM concurrence must be bounded by any specific bipartition, including the bipartition of the N qubits to party F and party G.

$$C_{GM}(|\Psi_i\rangle) \leq \sqrt{2}\sqrt{1 - \Pi_{F|G}(|\Psi_i\rangle)}. \tag{17}$$

Using the same mapping as done for Eq. 12 it is easy to show that $C(|\psi_i\rangle)$ is equal to the right hand side of Eq. 17. Therefore we conclude that

$$\begin{aligned}
C_{GM}(\hat{X}_{1i}) \leq \sum_i p_i C_{GM}(|\Psi_i\rangle) &\leq \sum_i p_i C(|\psi_i\rangle)\\
&= C(\hat{R}) = 2\max\{0, |z_1| - \sqrt{a_i b_i}\},
\end{aligned} \tag{18}$$

where the right most equality is found by evaluating Wootters's concurrence for $\hat{R}$ under the assumption $a_1b_1 \geq a_ib_i$. This upper bound matches the lower bound and therefore it is the exact value of $C_{GM}(\hat{X}_{1i})$. $\square$

Next, we generalize this result to all the X-matrices. We do so by decomposing the X-matrix into a convex sum of $\hat{X}_{1i}$ matrices. Let us first look at the case for which $|z_1| - w_1 \geq 0$.

(a) $|z_1| - w_1 \geq 0$. Note that $|z_i| - w_i \leq 0$ for $i \geq 2$ since $\sqrt{a_1b_1} \geq \sqrt{a_ib_i} \geq |z_i|$. First by a change of phase of the basis, which is a local unitary transformation, we change z_1 to $|z_1|$. This only changes the phase of the other off-diagonal elements. Then we decompose $\hat{X}$ in the following form.

$$\hat{X} = \hat{A} + \sum_{i>1}^{n} \hat{S}_i, \tag{19}$$

where $\hat{S}_i$ is an $\hat{X}_{1i}$ matrix whose two-qubit counterpart reads

$$\hat{R}_i = \begin{pmatrix} x_i & & & \sqrt{a_ib_i} \\ & a_i & z_i & \\ & z_i^* & b_i & \\ \sqrt{a_ib_i} & & & y_i \end{pmatrix}, \tag{20}$$

where

$$A_{11} = a_1\Big(1 - \frac{w_1}{\sqrt{a_1b_1}}\Big), \tag{21}$$

$$A_{2n,2n} = b_1\Big(1 - \frac{w_1}{\sqrt{a_1b_1}}\Big), \tag{22}$$

$$A_{1,2n} = A_{2n,1} = |z_1| - w_1, \tag{23}$$

$$A_{i,j} = 0 \quad i \neq \{1, 2n\}, \text{or} \quad j \neq \{1, 2n\}, \tag{24}$$

$$x_i = \frac{a_1\sqrt{a_ib_i}}{w_1}, \tag{25}$$

$$y_i = \frac{b_1\sqrt{a_ib_i}}{w_1}. \tag{26}$$

It can be shown that $\hat{S}_i$'s are all proportional to valid density matrices, since they are non-negative hermitian matrices. The proportionality constant is between zero and one and can be interpreted as probability. Using Lemma 1 one can show that all $\hat{S}_i$'s are biseparable matrices (though not normalized). Regarding the first matrix in the decomposition, the proportionality constant is $A_{11} + A_{2n,2n}$, and its GM concurrence is $2(|z_1| - w_1)/(A_{11} + A_{2n,2n})$. Due to the convexity of GM concurrence we conclude that

$$(A_{11} + A_{2n,2n})\frac{2(|z_1| - w_1)}{A_{11} + A_{2n,2n}} = 2(|z_1| - w_1) \tag{27}$$

is an upper bound for the GM concurrence of $\hat{X}$. Since $2(|z_1| - w_1)$ is also a lower bound for the concurrence, it is the exact value of the GM concurrence.

Note that for the above decomposition to work we had to assume that $|z_1| \geq w_1$. We now turn to the case $|z_1| < w_1$. We seek to show that all such density matrices are biseparable. We consider two different scenarios.

(b) $\sqrt{a_1b_1} \geq w_1$. In this case the matrix $\hat{X}$ can be decomposed to matrices similar to the previous case.

$$\hat{X} = \sum_{i>1}^{n} \hat{S}'_i, \tag{28}$$

where $\hat{S}'_i$ is an $\hat{X_{1i}}$ matrix whose two-qubit counterpart reads

$$\hat{R}'_i = \begin{pmatrix} a_1T_i & & & z_1T_i \\ & a_i & z_i & \\ & z_i^* & b_i & \\ z_1^*T_i & & & b_1T_i \end{pmatrix}, \quad T_i = \frac{\sqrt{a_ib_i}}{w_1}. \tag{29}$$

Since $|z_1|T_i \leq \sqrt{a_ib_i}$ and $|z_i| \leq \sqrt{a_1b_1}T_i$, we can invoke Lemma 1 to confirm $\hat{R}'_i$ is biseparable for all i. The fact that $\hat{R}'_i$ is not normalized does not interfere with the proof of biseparability as one can always factor out $\mathrm{Tr}[\hat{R}'_i]$. Now we focus on the last case.

(c) $\sqrt{a_1b_1} < w_1$. In this case we divide our matrix into two positive semi-definite matrices $\hat{X} = \hat{K_1} + \hat{L_1}$.

$$\hat{K_1} = \begin{pmatrix} a_1t & & & & & & & z_1t \\ & a_2r & & & & & z_2r & \\ & & \ddots & & & \iddots & & \\ & & & a_nr & z_nr & & & \\ & & & z_n^*r & b_nr & & & \\ & & \iddots & & & \ddots & & \\ & z_2^*r & & & & & b_2r & \\ z_1^*t & & & & & & & b_1t \end{pmatrix}, \tag{30}$$

$$\hat{L_1} = \begin{pmatrix} a_1r & & & & & & & z_1r \\ & a_2t & & & & & z_2t & \\ & & \ddots & & & \iddots & & \\ & & & a_nt & z_nt & & & \\ & & & z_n^*t & b_nt & & & \\ & & \iddots & & & \ddots & & \\ & z_2^*t & & & & & b_2t & \\ z_1^*r & & & & & & & b_1r \end{pmatrix}, \tag{31}$$

where

$$t = \frac{w_1}{w_1 + \sqrt{a_1b_1}}, \quad \text{and} \quad r = 1 - t.$$

Note that since $w_1 \leq 3\sqrt{a_1b_1}$, then $\frac{3}{4} \geq t > r$. One can show that

$$t\sqrt{a_1b_1} = rw_1, \tag{32}$$

which guarantees that $\hat{K_1}$ falls in the category of case (a). Since

$$\begin{aligned} &t|z_1| \leq rw_1, \\ &r|z_j| \leq (t - r)\sqrt{a_1b_1} + rw_j, \end{aligned} \tag{33}$$

$\hat{K_1}$ is biseparable. Regarding matrix $\hat{L_1}$, since

$$r|z_1| \leq tw_1,$$
$$t|z_j| \leq r\sqrt{a_1 b_1} + t\sum_{i\neq 1,j}^{n}\sqrt{a_i b_i}, \qquad (34)$$

it does not fall in the category of case (a) and thus belongs to either case (b) or case (c). If it falls in the category of case (b) then we can conclude that it is biseparable. If not, we divide $\hat{L_1}$ into two matrices $\hat{L_1} = \hat{K_2} + \hat{L_2}$, as before. Each time we divide a matrix in this way the trace of the remaining part is strictly smaller than the trace of the step before: $\mathrm{Tr}(\hat{L}_i) \leq 0.75^i$. Thus, we can write the matrix $\hat{X}$ as a convex sum of biseparable states and a remaining part that can be made arbitrarily close to zero. Therefore matrix $\hat{X}$ is a biseparable matrix. This completes the proof for all X-matrices. Therefore, we have proved that the GM concurrence of a N-qubit X-matrix is

$$C_{GM}(\hat{X}) = 2\max\{0, |z_i| - w_i\}, \quad 1 \leq i \leq n \qquad (35)$$

IV. N QUBITS IN N CAVITIES

The simple algebraic formula that was derived in the previous section opens up the possibility of analytically studying the entanglement dynamics of N qubits in different scenarios. The only restriction is that the density matrix of the system of interest remains an X-matrix. In this section we study the dynamics of the N-partite entanglement between N two-level atoms, each residing in its own single mode cavity, see Fig. 1. The interaction between each atom and its partner field is described by a Jaynes-Cummings Hamiltonian. The Hamiltonian for the total system reads

$$H_{tot} = \sum_i \frac{\omega_0}{2}\sigma_z^{(i)} + g(a_i^\dagger\sigma_-^{(i)} + a_i\sigma_+^{(i)}) + \omega a_i^\dagger a_i, \qquad (36)$$

where ω_0, ω are the frequencies of the atoms and cavities, respectively and g is the coupling constant between each atom and its partner cavity. For simplicity we have assumed that all the atoms and all of the cavities have the same resonance frequencies, ω_0 and ω, respectively. We assume that all of the atoms are initially in an N-qubit GHZ state, sharing N-partite entanglement, and the cavities are all in the vacuum state.

We present a detailed analysis only for the three qubit case but the generalization to N qubits is straightforward and we present the result for N qubits at the end. Our system is a generalization of the scenario previously studied by Yönaç *et al.* [22], so we follow the notation developed there. In fact, this generalization has already been employed in the study of multipartite entanglement in by Man *et al.* [25], where the entanglement of different biseparations of the system were studied. Although separability of a particular bipartition of the system is a sufficient condition for biseparability it is not a necessary condition. Thus a quantitative examination of N-partite entanglement was missing; this is the analysis that we seek to provide using our formula. For simplicity from now on we refer to GM concurrence simply as concurrence.



FIG. 1. A schematic representation of the setup where the remote atoms are interacting with their own cavities. There is no interaction between remote parties. The qubits are assumed to be initially maximally entangled and the fields are in individual separable vacuum states.

We first consider an initial state which is an analog of the so-called $|\Phi,\alpha\rangle$ states for three atoms [22] . The initial state of the system reads

$$|\Phi,\alpha\rangle = (\cos\alpha|e,e,e\rangle + \sin\alpha|g,g,g\rangle) \otimes |0,0,0\rangle. \qquad (37)$$

where $|e\rangle$ denotes the excited state of a two level atom $\sigma_z|e\rangle = |e\rangle$ and $|g\rangle$ denotes the ground state $\sigma_z|g\rangle = -|g\rangle$. All three cavities are assumed to be in their ground states initially. Here we use a slightly different notation than we used in the previous section, where 0 and 1 were used for bits of a qubit. From now on, 0 and 1 will be used for the vacuum state and a singly excited state of the cavity. For each atom and cavity the two states $|e,0\rangle$ and $|g,0\rangle$ will evolve according to

$$U(t)|e,0\rangle = A_t|e,0\rangle + B_t|g,1\rangle,$$
$$U(t)|g,0\rangle = |g,0\rangle, \qquad (38)$$

where $U(t)$ is the Jaynes-Cummings propagator. If the atoms are in resonance with the cavities ($\omega_0 = \omega$), then one can show that

$$A_t = e^{i\omega t}\cos gt, \qquad B_t = -ie^{-i\omega t}\sin gt. \qquad (39)$$

Thus our initial state evolves to

$$\begin{aligned}|\Phi(t),\alpha\rangle = \cos\alpha\ &(A_t|e,0\rangle + B_t|g,1\rangle)\\ &\otimes(A_t|e,0\rangle + B_t|g,1\rangle)\\ &\otimes(A_t|e,0\rangle + B_t|g,1\rangle)\\ &+\sin\alpha\ |g,0\rangle\otimes|g,0\rangle\otimes|g,0\rangle. \qquad (40)\end{aligned}$$

By tracing out the cavities we find the density matrix of



FIG. 2. $Q_N(t)$ for different number of qubits. $N = 2$ (dashed line), $N = 10$ (dottedd line), $N = 20$ (dashed-dotted line), and $N = 100$ (solid line). The initial states are $|\Phi_N, \frac{\pi}{4}\rangle$.



FIG. 3. $R_N(t)$ and $T_N(t)$ for different number of qubits. $N = 2$ (dashed line), $N = 10$ (dottedd line), $N = 20$ (dashed-dotted line), and $N = 100$ (solid line). The initial states are $|\Phi_N, \frac{\pi}{4}\rangle$.

the three atoms,

$$\hat{\rho}_{\Phi_3}(t) = \begin{pmatrix} a_1 & & & & & & & z_1 \\ & a_2 & & & & & & \\ & & a_2 & & & & & \\ & & & b_2 & & & & \\ & & & & a_2 & & & \\ & & & & & b_2 & & \\ & & & & & & b_2 & \\ z_1 & & & & & & & b_1 \end{pmatrix}, \tag{41}$$

where

$$\begin{aligned} a_1 &= \cos^2\alpha |A_t|^6, \\ b_{1=} &\sin^2\alpha + \cos^2\alpha |B_t|^6, \\ z_1 &= \sin\alpha\cos\alpha A_t^3, \\ a_2 &= \cos^2\alpha |A_t^2 B_t|^2, \\ b_2 &= \cos^2\alpha |A_t B_t^2|^2. \end{aligned} \tag{42}$$

The concurrence of this density matrix is

$$\begin{aligned} &C_3(t) = \max\{0, Q_3(t)\}, \\ &Q_3(t) = 2\cos^2\alpha |\cos^3 gt| \left(|\tan\alpha| - 3|\sin^3 gt| \right). \end{aligned} \tag{43}$$

If, instead of the three qubit-cavity systems, we had N qubit-cavities then one can show that the concurrence of the N qubits would be

$$\begin{aligned} &C_N(t) = \max\{0, Q_N(t)\}, \\ &Q_N(t) = R_N(t)\; T_N(t), \\ &R_N(t) = 2\cos^2\alpha |\cos^N gt|, \\ &T_N(t) = |\tan\alpha| - (2^{N-1} - 1)|\sin^N gt|. \end{aligned} \tag{44}$$

For $N = 2$, we recover the formula given by Yönaç *et al.* [22]. We plot Q_N, T_N and R_N in Fig. 2 and Fig. 3 for different numbers of qubits ranging from 2 to 100. As N increases, the positive part of T_N approaches a step function (when $T_N \leq 0$, the concurence is zero), whereas R_N remains non-negative and only vanishes momentarily. Thus T_N determines when entanglement vanishes completely and R_N determines the magnitude of concurrence. Notably $R_N(t) = R_{N-1}|\cos gt|$, which guarantees that a as N increases the bulk of the initial concurrence decays faster. The time it takes for the concurrence to decay by a factor of two can be approximated by

$$gT_{\frac{1}{2}} = \arccos\frac{1}{\sqrt[N]{2}}. \tag{45}$$

The other time of interest is when the concurrence vanishes completely. For $N \geq 3$, we can directly confirm the phenomenon of multipartite entanglement sudden death (ESD). To find the time of ESD, T_{ESD}, we solve the equation $Q_N = 0$ which gives

$$\begin{aligned} T_{ESD} &= \frac{1}{g}\arcsin\left|\frac{\tan\alpha}{2^{N-1}-1}\right|^{\frac{1}{N}}, \\ \lim_{N\to\infty} T_{ESD} &= \frac{\pi}{6g}. \end{aligned} \tag{46}$$

In the limit of large N, the time of ESD is independent of the initial concurrence; it depends only on the common coupling strength g. In Fig. 4 we have plotted the ESD time against the number of qubits. When the initial amount of concurrence is considerable the ESD time becomes smaller as N increases. However, when the initial concurrence is small the T_{ESD} increases and converges to the limit $T_{ESD} = \pi/6g$ as N tends to infinity. It is notable that even for a very large N, although the bulk on initial entanglement decays very rapidly, the state of the two-level atoms remains entangled and becomes biseparable only after T_{ESD}.

We also consider an initial state which is the three-qubit analog of the so-called $|\Psi\rangle$ state.

$$|\Psi, \alpha\rangle = (\cos\alpha|e, e, g\rangle + \sin\alpha|g, g, e\rangle) \otimes |0, 0, 0\rangle \tag{47}$$

This state evolves according to

$$\begin{aligned} |\Psi(t), \alpha\rangle = &\cos\alpha\; (A_t|e, 0\rangle + B_t|g, 1\rangle) \\ &\otimes (A_t|e, 0\rangle + B_t|g, 1\rangle) \otimes |g, 0\rangle \\ &+ \sin\alpha\; |g, 0\rangle \otimes |g, 0\rangle \otimes (A_t|e, 0\rangle + B_t|g, 1\rangle). \end{aligned} \tag{48}$$



FIG. 4. ESD time versus number of qubits for different initial states $|\Phi_N, \alpha\rangle$. From the bottom up $\tan\alpha = 0.01, 0.1, 0.2, 0.5$, and 1 respectively.



FIG. 5. $Q_N(t)$ versus time for different numbers of qubits. $N = 2$ (dashed line), $N = 10$ (dottedd line), $N = 20$ (dashed-dotted line), and $N = 100$ (solid line). The initial states are a $|\Psi_N, \frac{\pi}{4}\rangle$

The reduced density matrix of the three atoms in the basis $\{|e,e,e\rangle, |e,e,g\rangle, |e,g,e\rangle, \ldots, |g,g,g\rangle\}$ is

$$\rho_{\Psi_3}(t) = \begin{pmatrix} 0 & & & & & & & \\ & a_2 & & & & & z_2 & \\ & & 0 & & & & & \\ & & & a_4 & & & & \\ & & & & 0 & & & \\ & & & & & b_3 & & \\ & z_2 & & & & & b_2 & \\ & & & & & & & b_1 \end{pmatrix} \tag{49}$$

where

$$\begin{aligned} a_2 &= \cos^2\alpha |A_t|^4 \\ z_2 &= \sin\alpha\cos\alpha A_t^2 A_t^* \\ a_4 &= \cos^2\alpha |A_t B_t|^2 \\ b_2 &= \sin^2\alpha |A_t|^2 \\ b_3 &= \cos^2\alpha |A_t B_t|^2 \\ b_1 &= \sin^2\alpha |B_t|^2 + \cos^2\alpha |B_t|^4 \end{aligned} \tag{50}$$

The concurrence is given by

$$C_3(t) = 2|\sin\alpha\cos\alpha\cos^3 gt|. \tag{51}$$

This state does not experience ESD due to the fact that $\langle e,e,e|\rho_\Psi|e,e,e\rangle$, $\langle e,g,e|\rho_\Psi|e,g,e\rangle$, and $\langle g,e,e|\rho_\Psi|g,e,e\rangle$ remain zero for all times. To show this, we note that our initial state is a superposition of two possibilities. Either two of the atoms, $\{1,2\}$, are excited and the other atom, $\{3\}$, is in vacuum state, or that single atom is excited and the atoms $\{1,2\}$ are in their ground states. Since all cavities are initially in their ground states, if an atom is in its ground state initially it will always remain there. But the three diagonal terms that we referred to require the atom $\{3\}$ and at least one of the other two atoms be excited simultaneously. Since this is forbidden, all of these matrix elements remain zero. Thus $\sqrt{a_1 b_1} = \sqrt{a_3 b_3} = \sqrt{a_4 b_4} = 0$, and $C_3 = 2|z_2|$. This argument can be generalized for $N \geq 4$ to all of the GHZ states except for $|\Phi_N, \alpha\rangle$, because in the $|\Phi_N, \alpha\rangle$ state all of the atoms can be initially excited. Therefore none of the GHZ states except $|\Phi_N, \alpha\rangle$ experience ESD. The concurrence of all such GHZ states is given by

$$C_N(t) = 2|\sin\alpha\cos\alpha\cos^N gt|. \tag{52}$$

In Fig. 5 we plot Q_N for different numbers of qubits ranging from 2 to 100, The initial state is $|\Psi_N, \frac{\pi}{4}\rangle$. As expected from the concurrence formula, ESD does not happen for any N and only vanishes momentarily. Still, as N increases the width of the periods of considerable concurrence decreases.

V. CONCLUSION

We have found an algebraic formula for the genuinely multipartite (GM) concurrence of N-qubit density matrices that can be written as X-matrices in an orthonormal product basis. This development allows N-partite entanglement to be quantified for such states. The formula opens up the possibility of studying entanglement dynamics of N-qubit states in different scenarios, as long as the X-form of the density matrix is preserved. This includes scenarios such as local decoherence channels that will be studied elsewhere.

Using the concurrence formula, we have studied the dynamics of N-partite entanglement of N two-level atoms interacting with partner single-mode cavities. We showed that the so-called $|\Phi_N, \alpha\rangle$ states experience ESD whereas the $|\Psi_N, \alpha\rangle$ states do not experience ESD. Algebraic formulas for the concurrence of the two different case were presented. Since the entanglement dynamics is periodic we focused on how fast the initial entanglement vanishes in the initial cycle before reviving to its initial value at $gt = 2\pi$. It is observed that although the bulk of initial concurrence decays faster as the N increases, the same conclusion can not be drawn about the exact moment at which the entanglement disappears, T_{ESD}. In fact for the states that have small initial entanglement, T_{ESD} is prolonged as the number of qubits increases.

VI. ACKNOWLEDGEMENT

S.M.H.R. acknowledges useful communication with O. Gühne. We acknowledge partial financial support from ARO W911NF-09-1-0385 and NSF PHY-0855701. M.H. gratefully acknowledges support from the EC-project IP "Q-Essence", the ERC advanced grant "IRQUAT" and MC grant "Quacocos".

---

[1] E. Schrödinger, Proc. Cambridge Philos. Soc. **31**, 555 (1935).

[2] E. Schmidt, Math. Annalen **63**, 433 (1906).

[3] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Rev. Mod. Phys. **81**, 865 (2009).

[4] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quanum Information* (Cambridge University Press, 2000).

[5] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).

[6] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996); M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Lett. A **223**, 1 (1996).

[7] B. M. Terhal and K. G. H. Vollbrecht, Phys. Rev. Lett. **85**, 2625 (2000); P. Rungta and C. M. Caves, Phys. Rev. A **67**, 012307 (2003); R. Lohmayer, A. Osterloh, J. Siewert, and A. Uhlmann, Phys. Rev. Lett. **97**, 260502 (2006); T.-C. Wei, Phys. Rev. A **78**, 012327 (2008).

[8] V. Giovannetti, S. Lloyd, and L. Maccone, Science **306**, 1330 (2004).

[9] T. R. de Oliveira, G. Rigolin, and M. C. de Oliveira, Phys. Rev. A **73**, 010305 (2006); L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Rev. Mod. Phys. **80**, 517 (2008).

[10] W. Dür and H.-J. Briegel, Phys. Rev. Lett. **92**, 180403 (2004).

[11] A. R. R. Carvalho, F. Mintert, and A. Buchleitner, Phys. Rev. Lett. **93**, 230501 (2004).

[12] C. Simon and J. Kempe, Phys. Rev. A **65**, 052327 (2002).

[13] L. Aolita, R. Chaves, D. Cavalcanti, A. Acín, and L. Davidovich, Phys. Rev. Lett. **100**, 080501 (2008).

[14] O. Gühne and M. Seevinck, New Journal of Physics **12**, 053002 (2010).

[15] B. Jungnitsch, T. Moroder, and O. Gühne, Phys. Rev. Lett. **106**, 190502 (2011).

[16] S. Szalay, Phys. Rev. A **83**, 062337 (2011).

[17] Z.-H. Ma, Z.-H. Chen, J.-L. Chen, C. Spengler, A. Gabriel, and M. Huber, Phys. Rev. A **83**, 062325 (2011).

[18] D. T. Pope and G. J. Milburn, Phys. Rev. A **67**, 052107 (2003).

[19] P. Love, A. van den Brink, A. Smirnov, M. Amin, M. Grajcar, E. Ilichev, A. Izmalkov, and A. Zagoskin, Quant. Inf. Proc. **6**, 187 (2007).

[20] P. Rungta, V. Bužek, C. M. Caves, M. Hillery, and G. J. Milburn, Phys. Rev. A **64**, 042315 (2001).

[21] T. Yu and J. H. Eberly, Quantum Inf. Comput. **7**, 459 (2007).

[22] M. Yönaç, T. Yu, and J. H. Eberly, J. Phys. B: At. Mol. Opt. Phys. **39**, S621 (2006).

[23] T. Yu and J. H. Eberly, Phys. Rev. Lett. **93**, 140404 (2004).

[24] M. Yönaç, T. Yu, and J. H. Eberly, J. Phys. B: At. Mol. Opt. Phys. **40**, S45 (2007); H. T. Cui, K. Li, and X. X. Yi, Phys. Lett. A **365**, 44 (2007); G. Zhang and Z. Chen, Opt. Commun. **275**, 274 (2007); Z. Ficek and R. Tanaś, Phys. Rev. A **77**, 054301 (2008); A. Al-Qasimi and D. F. V. James, *ibid.* **77**, 012117 (2008); M. Ali, G. Alber, and A. R. P. Rau, Journal of Physics B: At. Mol. Opt. Phys. **42**, 025501 (2009).

[25] Z.-X. Man, Y.-J. Xia, and N. B. An, Journal of Physics B: At. Mol. Opt. Phys. **41**, 155501 (2008).

[26] Y. S. Weinstein, Phys. Rev. A **82**, 032326 (2010); L. Aolita, D. Cavalcanti, A. Acín, A. Salles, M. Tiersch, A. Buchleitner, and F. de Melo, *ibid.* **79**, 032322 (2009); Y. S. Weinstein, *ibid.* **79**, 012318 (2009).

[27] O. Gühne, F. Bodoky, and M. Blaauboer, Phys. Rev. A **78**, 060301 (2008).

[28] M. Huber, F. Mintert, A. Gabriel, and B. C. Hiesmayr, Phys. Rev. Lett. **104**, 210501 (2010).